

Effective Theory Description of Weak Annihilation in $B \rightarrow X_u \ell \nu$ Decays

J.J. Sanz-Cillero^{a*}

^a Groupe Physique Théorique, IPN Orsay, Université Paris-Sud XI, 91406 Orsay, France
e-mail: cillero@ipno.in2p3.fr

The semileptonic $B \rightarrow X_u \ell \nu$ decays allow a pretty clean determination of the CKM matrix element $|V_{ub}|$. Nevertheless, the presence of weak annihilation effects near the end-point of the q^2 spectrum introduces uncertainties in the inclusive calculation, requiring the use of non-perturbative techniques like heavy meson chiral perturbation theory and large N_C limit.

1. Introduction

The $B_q \sim b\bar{q}$ meson decay through the inclusive semileptonic channel $B_q \rightarrow X_u \ell \nu$ has been shown to be a very clean way to determine $|V_{ub}|$ [2].

The inclusive B_q analysis makes use of the operator product expansion (OPE) [2,3]. The first difference between the B_u^- and $B_{d,s}^0$ decays arises at next-to-next-to-next-to-leading order (N³LO) in the $1/m_b$ expansion. It is due to the topologies where the incoming $b\bar{q}$ pair annihilates into a W -boson (Fig. (1)), giving away an infinitesimal amount of momentum through soft gluon radiation. These contributions are known as weak-annihilation (WA) [4]. They appear as a delta function at the end-point of the spectrum $q^2 = (p_\ell + p_\nu)^2 = m_b^2$. Their determination is rather relevant if the uncertainty on $|V_{ub}|$ is to be reduced up to a few per cent [2].

In order to better isolate the WA, the analysis is carried on within the chiral limit $m_q = 0$.

2. OPE and Weak-Annihilation

At $\mathcal{O}(1/m_b^3)$, the WA in the OPE is given in terms of the bag constants $B_{WA} \equiv B_2 - B_1$ [5]:

$$\Gamma(B_q \rightarrow X_u \ell \nu) = \dots + \frac{G_F^2 |V_{ub}|^2}{12\pi} f_B^2 m_b^3 B_{WA}^q, \quad (1)$$

*Thanks to the organisers of BEACH 2006, Lancaster, UK, and to C. Smith, E. Gardi, M.Ciuchini and L. Oliver for useful comments and suggestions. Also thanks to M. Neubert for his help and contributions in early stages of the project. Work done in collaboration with G. Paz [1]. and supported by EU RTN Contract CT2002-0311.




Figure 1. Examples of a) non-annihilating topologies; b) WA diagrams.

with $B_{WA}^q = 0$ in the factorisation limit, independently of the light quark in the B_q meson ($B_2^{d,s} = B_1^{d,s} = 1$ and $B_2^u = B_1^u = 0$). The expected 10% deviations from the factorisation limit provide an estimate of the WA contribution to the branching ratio (BR) [5]:

$$\delta\mathcal{B}_{B_q \rightarrow X_u \ell \nu} = 3.9 \times \left(\frac{f_B}{0.2 \text{ GeV}} |V_{ub}| \right)^2 \left(\frac{B_{WA}^q}{0.1} \right). \quad (2)$$

Other theoretical and experimental estimates provide similar values [6,7].

The OPE calculation also points out relevant aspects about the nature and the scalings of the WA contributions:

- Constant contribution to the width, generated at the end-point, in the non-perturbative region of low hadronic energies $E_X \lesssim 1 \text{ GeV}$ [4].

- The WA operators appear at N³LO in the heavy mass expansion, so their contributions to

the BR is $\mathcal{O}(\Lambda_{\text{QCD}}^3/m_b^3)$.

–The WA topologies introduce an extra quark loop in the hadronic tensor $T^{\mu\nu}$, being suppressed by $1/N_C$ in the large N_C limit [8].

–The WA $B^- - B^0$ difference is related to the production of chiral singlets in the final state X_u (see Fig. (1)). In the chiral limit, WA is the responsible of the difference between the B^+ and the B^0 widths.

–At N³LO in $1/m_b$ there are other non-WA terms in the OPE (independent of q in B_q) that need to be taken into account and properly disentangled [6].

3. Cut-off on the hadronic energy E_X

In order to isolate the WA effects and remove the $B \rightarrow X_c \ell \nu$ background we consider the hadronic energy cut-off $E_X < \Lambda_E$.

Multi-pion production is found to be suppressed both experimentally [9] and theoretically. The analysis of the $B \rightarrow \pi \pi \ell \nu$ channel through Heavy Meson Chiral perturbation Theory (HM χ PT) [10] finds it widely suppressed with respect to $B \rightarrow \pi \ell \nu$ for $\Lambda_E \lesssim 1$ GeV [1].

Hence, only the lightest pseudo-scalar meson production $B \rightarrow P \ell \nu$ is considered here. The decay into light vector mesons $B \rightarrow V \ell \nu$ introduces small corrections for $\Lambda_E \lesssim 1$ GeV and it can be neglected in first approximation [1].

In the $B_q \rightarrow P \ell \nu$ mode, one has $P_X^2 = m_P^2$ and $E_X = E_P$, with the phase-space given by

$$\frac{1}{2} \left(z - \sqrt{z^2 - 4r^2} \right) \leq \bar{x} \leq \frac{1}{2} \left(z + \sqrt{z^2 - 4r^2} \right), \quad 2r \leq z \leq \epsilon, \quad (3)$$

where $z = \frac{2E_P}{M_B}$, $\bar{x} = 1 - \frac{2E_\ell}{M_B}$, $\epsilon = \frac{2\Lambda_E}{M_B}$ and $r = \frac{m_P}{M_B}$.

The decay is determined by the matrix element $\langle P | \bar{u} \gamma^\mu (1 - \gamma_5) b | B \rangle = (p_B + p_P)^\mu f_+ + q^\mu f_-$, (4) although, for $m_\ell = m_\nu = 0$, only the form factor $f_+(z)$ contributes to the width:

$$\frac{1}{\Gamma_0} \frac{d\Gamma}{dz} = |f_+(z)|^2 \cdot [z^2 - 4r^2]^{\frac{3}{2}}. \quad (5)$$

with $\Gamma_0 = \frac{G_F^2 |V_{ub}|^2 M_B^5}{192\pi^3}$. Integrating the energy on the range $2r \leq z \leq \epsilon$ provides the corresponding contribution to the width $\Delta\Gamma(\epsilon, r_\epsilon)$ from this part

of the phase-space. It only depends on ϵ and the ratio $r_\epsilon = \frac{2r}{\epsilon} = \frac{m}{\Lambda_E}$.

4. HM χ PT and large N_C limit

To compute $f_+(z)$ and $\Delta\Gamma(\epsilon, r_\epsilon)$ we make use of HM χ PT. At large N_C , the meson loops are suppressed. The $B \rightarrow P \ell \nu$ amplitude is dominated the B^* pole from the tree-level exchange. Its description is provided by the leading HM χ PT lagrangian [10]:

$$|f_+(z)|_{B_q \rightarrow P}^2 = D_{B_q \rightarrow P} \cdot \frac{f_B^2 g^2}{2 F_\pi^2} \frac{1}{z^2}, \quad (6)$$

with the group factors $D_{B_d \rightarrow \pi^+} = D_{B_s \rightarrow K^+} = 1$, $D_{B_u \rightarrow \pi^0} = \frac{1}{2}$, $D_{B_u \rightarrow \eta_8} = \frac{1}{6}$, $D_{B_u \rightarrow \eta_0} = \frac{1}{3}$. f_B and $F_\pi \simeq 92.4$ MeV are the B and π decay constants, respectively, and g is the $B\pi B^*$ coupling [10].

In the large N_C and chiral limits, the pseudoscalars (π, K, η_8, η_0) are massless, so $r_\epsilon = 0$. Therefore, the integrated widths coincide:

$$\Delta\Gamma(\epsilon)_{B_d^0, B_s^0, B_u^+ \rightarrow X_u \ell \nu} = \frac{f_B^2 g^2}{4 F_\pi^2} \cdot \epsilon^2, \quad (7)$$

given by the decays $B_d^0 \rightarrow \pi^+$, $B_s^0 \rightarrow K^+$ and $B_u^- \rightarrow \pi^0, \eta_8, \eta_0$. Notice that at large N_C the η_0 forms the ninth chiral Goldstone boson.

The modifications to the interaction vertices and couplings in the form factors (Eq. (6)) are expected to be of the order of $\frac{1}{N_C}$ and $\frac{\Lambda_{\text{QCD}}}{m_b}$.

However, the gaining of mass by the η_0 meson is far more important. It generates large modifications in the available phase-space in Eq. (3) and important kinematical corrections, of the order of $\left(\frac{m_{\eta_0}}{\Lambda_E}\right)^2$. The chiral octet (π, K, η_8) remains massless within the chiral limit whereas the chiral singlet η_0 gains a mass $m_{\eta_0} \sim 900$ MeV due to $1/N_C$ corrections [11]. The available phase-space for the $B_u^- \rightarrow \eta_0$ mode gets reduced and the integration of the form factor (Eq. (6)) over the available range of z yields

$$\Delta\Gamma(\epsilon, r_\epsilon)_{B_u \rightarrow \eta_0} = \frac{1}{3} \times \frac{f_B^2 g^2}{4 F_\pi^2} \times \epsilon^2 \times \left[(1 + 2r_\epsilon^2) \sqrt{1 - r_\epsilon^2} + \frac{3r_\epsilon^2}{2} \ln \frac{1 + \sqrt{1 - r_\epsilon^2}}{1 - \sqrt{1 - r_\epsilon^2}} \right]. \quad (8)$$

The other decay modes remain unaffected.

5. Results and conclusions

By observing the $B^0 - B^-$ difference it is possible to separate the WA from other $1/m_b^3$ effects [6]. The subtraction of the integrated widths $\Delta\Gamma(\epsilon)_{B_q \rightarrow X_u \ell \nu}$ gives

$$\frac{[\Delta\Gamma_{B^-} - \Delta\Gamma_{B^0}]}{\Gamma_0} = \frac{f_B^2 g^2 m_{\eta_0}^2}{2F_\pi^2 M_B^2} \left\{ 1 + \ln \frac{r_\epsilon^2}{4} \right\} + \dots (9)$$

with the dots standing for contributions suppressed by $\left(\frac{m_{\eta_0}}{\Lambda_E}\right)^2$. This result reproduces all the features in the difference $[\Delta\Gamma_{B^-} - \Delta\Gamma_{B^0}]$ that one expects from the OPE:

- Since $f_B^2 \sim 1/M_B$, suppression by $1/M_B^3$.
- Suppression by $1/N_C$: from the scaling $m_{\eta_0}^2 \sim \frac{1}{N_C}$ with respect to the other energy scales.
- The production of chiral singlets in the final state is responsible of the difference (WA topologies in Fig. (1)).

- As Λ_E grows, the difference $\Delta\Gamma_{B^-} - \Delta\Gamma_{B^0}$ tends to a constant up to a residual term $\ln\left(\frac{m_{\eta_0}}{\Lambda_E}\right)$; the effect is non-perturbative and comes from the end-point of the q^2 spectrum. The removal of this residual log needs the consideration of higher B multiplets, in order to recover the $f_+ \sim \frac{1}{E_P^2}$ behaviour of the form factor at high energies [12]. This have been implemented by considering modified-pole expressions [13].

In Fig. (2), we show the dependence of the difference of branching ratios $[\mathcal{B}_{B_d^0} - \mathcal{B}_{B_u^-}]$ on the cut-off, ϵ . One can see how around $\Lambda_E \sim 1$ GeV, the difference stabilises and becomes approximately constant (upto the residual log dependence referred to before), reaching the value

$$[\Delta\mathcal{B}_{B_d^0} - \Delta\mathcal{B}_{B_u^-}] \sim 4 \times \left(\frac{f_B}{0.2 \text{ GeV}} \frac{g}{0.5} |V_{ub}| \right)^2, (10)$$

in agreement with former estimates [5,6,7].

This WA estimate shows how it is possible to describe these OPE effects through HM χ PT, recovering all the expected features.

The next step is the calculation of the $B \rightarrow V \ell \nu$ contribution. The residual $\ln(r_\epsilon)$ dependence in Eq. (9) is expected to be removed through modified pole expressions [13]. The relevance of this effect and the proposed techniques should be taken into account in the analyses of D decays, where WA will be more important.




Figure 2. Dependence of the difference of branching ratios $[\Delta\mathcal{B}_{B_d^0} - \Delta\mathcal{B}_{B_u^-}]$ on the cut-off ϵ (dashed curve). For comparison, it is shown together with $\Delta\mathcal{B}_{B_d^0}$ (solid). Expressions are from Eqs. (7)–(9) and $\Delta\mathcal{B}$ is expressed in units of $\left(\frac{f_B}{0.2 \text{ GeV}} \frac{g}{0.5} |V_{ub}|\right)^2$.

REFERENCES

1. G. Paz and J.J. Sanz-Cillero; in preparation.
2. B.O. Lange, M. Neubert and G. Paz, *Phys. Rev. D* **72** (2005) 073006.
3. F. De Fazio and M. Neubert, *JHEP* **06** (1999) 017; M. Neubert and T. Becher, *Phys. Lett. B* **535** (2002) 127.
4. I.I.Y. Bigi and N.G. Uraltsev, *Nucl. Phys. B* **423** (1994) 33-55.
5. M.B. Voloshin, *Phys. Lett. B* **515** (2001) 74-80.
6. P. Gambino, G. Ossola and N. Uraltsev, *JHEP* **09** (2005) 010.
7. T.O. Meyer, Ph.D. Thesis, Cornell U. (2005).
8. G. 't Hooft, *Nucl. Phys. B* **72** (1974) 461; *Nucl. Phys. B* **75** (1974) 461; E. Witten, *Nucl. Phys. B* **160** (1979) 57.
9. Particle Data Group (S. Eidelman et al.); *Phys. Lett. B* **592** (2004) 1.
10. G. Burdman and H.F. Donoghue, *Phys. Lett. B* **280** (1992) 287-291; M.B. Wise, *Phys. Rev. D* **45** (1992) 2188-2191.
11. R. Kaiser, hep-ph/0502065; J.M. Gérard and E. Kou, *Phys. Lett. B* **616** (2005) 85-92; J.M. Gérard, C. Smith and S. Trine, *Nucl. Phys. B* **730** (2005) 1-36.
12. J. Charles *et al.*, *Phys. Rev. D* **60** (1999) 014001.
13. D. Becirevic and J. Kaidalov, *Phys. Lett. B* **478** (2000) 417; P. Ball and R. Zwicky, *Phys. Lett. B* **625** (2005) 225; S. Fajfer and J. Kamenik, *Phys. Rev. D* **71** (2005) 014020.